Page: 36 Run Date: 12/31/03

| Race/Ethnicity and Year | | | Smo | king | | Quit Smoking By First Prenatal | | Smoking in Household | | | |
|-------------------------|----------------------|--------------------------------|----------------------|-------------------------------|----------------------|--|----------------------|-------------------------------|----------------------|-------------------------------|----------------------|
| | | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | Visit and Stayed Off Cigarettes (3) | | Prenatal | | Postpartum | |
| | | Number | % | Number | % | Number | % | Number | % | Number | % |
| White, Not Hispanic | 2002 2001 2000 | 304,300 309,340 298,682 | 47.3 48.0 47.8 | 232,827 234,066 232,242 | 30.1 31.2 31.2 | 85,806 84,905 84,680 | 28.6 27.3 27.1 | 167,068 166,389 147,659 | 39.1 40.0 41.2 | 186,646 188,356 168,950 | 34.8 36.2 37.4 |
| Black, Not Hispanic | 2002 2001 2000 | 137,123 139,875 139,192 | 17.9 18.5 18.6 | 92,458 91,355 89,395 | 12.4 13.1 13.3 | 14,289 14,075 13,930 | 37.0 35.1 33.9 | 58,705 61,159 58,003 | 17.8 18.0 18.9 | 63,772 64,682 60,260 | 13.8 14.2 15.1 |

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Excludes data with errors.

⁽³⁾ Among women who smoked 3 months prior to pregnancy.

^{*} Percentages are not calculated if <100 records are available for analysis after exclusions.

Page: Run Date: 12/31/03

37

| Summary of Trends in Smoking Indicators by Race/Ethnicity (2) | | | | | | | | | | | | |
|---|----------------------|--------------------------------|----------------------|----------------------------|----------------------|--|----------------------|----------------------------|----------------------|----------------------------|----------------------|--|
| Race/Ethnicity and Year | | Smoking | | | | Quit Smoking By First Prenatal | | Smoking in Household | | | | |
| | | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | Visit and Stayed Off Cigarettes (3) | | Prenatal | | Postpartum | | |
| | | Number | % | Number | % | Number | % | Number | % | Number | % | |
| Hispanic | 2002 2001 2000 | 118,957 114,020 102,408 | 6.0 6.5 6.8 | 71,267 64,575 57,576 | 2.6 2.8 3.3 | 3,448 3,208 3,021 | 54.1 55.5 51.8 | 59,219 57,556 50,596 | 7.4 8.0 8.3 | 65,539 61,370 53,317 | 6.0 6.4 7.0 | |
| American Indian/Alaskan Native | 2002 2001 2000 | 7,110 6,794 6,073 | 37.8 39.7 36.9 | 6,362 5,955 5,343 | 18.4 20.0 20.6 | 1,696 1,681 1,412 | 36.8 35.2 33.9 | 4,892 4,172 3,394 | 28.7 31.0 28.6 | 4,966 4,468 3,514 | 24.5 26.9 24.8 | |

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Excludes data with errors.

⁽³⁾ Among women who smoked 3 months prior to pregnancy.

^{*} Percentages are not calculated if <100 records are available for analysis after exclusions.

Page: 38 Run Date: 12/31/03

| | Sur | nmary o | f Trends | s in Smok | ing Indi | cators by | Race/E | thnicity | (2) | | | |
|-------------------------|----------------------|--------------------------------|----------------------|----------------------------|--------------------|--|----------------------|--------------------------|----------------------|----------------------------|----------------------|--|
| Race/Ethnicity and Year | | | Smo | oking | | Quit Smoking By First Prenatal | | Smoking in Household | | | | |
| | | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | Visit and Stayed Off Cigarettes (3) | | Prenatal | | Postpartum | | |
| | | Number | % | Number | % | Number | % | Number | % | Number | % | |
| Asian/Pacific Islander | 2002 2001 2000 | 15,445 15,385 15,464 | 12.8 13.1 10.5 | 14,760 14,348 13,815 | 4.9 5.4 5.1 | 1,480 1,410 1,172 | 56.0 56.6 50.3 | 10,126 9,983 9,521 | 21.8 20.5 15.7 | 13,123 12,461 11,380 | 13.9 17.7 14.9 | |
| All Other/Unknown | 2002 2001 2000 | 8,169 7,828 6,428 | 19.0 15.7 14.5 | 6,081 5,570 4,391 | 11.7 8.8 8.8 | 809 623 463 | 34.0 37.7 34.6 | 5,682 5,840 4,373 | 17.2 15.8 15.1 | 6,854 6,452 4,837 | 16.5 12.6 12.3 | |

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Excludes data with errors.

⁽³⁾ Among women who smoked 3 months prior to pregnancy.

^{*} Percentages are not calculated if <100 records are available for analysis after exclusions.

Page:

Run Date: 12/31/03

| Summary of Trends in Smoking Indicators by Race/Ethnicity (2) | | | | | | | | | | | | | |
|---|----------------------|-------------------------------|--------------------------------|-------------------------------|----------------------------|-----------------------------------|--|-------------------------------|----------------------|-------------------------------|----------------------|--|--|
| Race/Ethnicity and Year | | Smoking | | | | Quit Smoking By First Prenatal | | Smoking in Household | | | | | |
| | | | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | Visit and Stayed Off Cigarettes (3) | | Prenatal | | artum | | |
| | | Number | % | Number | % | Number | % | Number | % | Number | % | | |
| | | | | | | | | | | | | | |
| Total All Race/Ethnic Groups | 2002 2001 2000 | 591,104 593,242 568,247 | 30.8 31.6 31.7 | 423,755 415,869 402,762 | 20.3 21.5 22.0 | 107,528 105,902 104,678 | 31.1 29.7 29.1 | 305,692 305,099 273,546 | 27.7 28.4 28.9 | 340,900 337,789 302,258 | 24.0 25.3 26.2 | | |
| | | | | | | | | | | | | | |
| | | | | | | | | | | | | | |



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Excludes data with errors.

⁽³⁾ Among women who smoked 3 months prior to pregnancy.

Percentages are not calculated if <100 records are available for analysis after exclusions.